Problem sheet 2: Landau levels and quantum Hall effect **20.10.2026**

2.1. Cyclotron motion

Consider a classical particle of mass m and charge $q = -e$ moving in two dimensions in the presence of a perpendicular magnetic field of magnitude $\mathbf{B} = B\mathbf{e}_z$, so that this particle executes the standard 2D cyclotron motion. Answer the following questions about the cyclotron motion:

(a) Consider the two cases, 1 and 2, in which the particle is prepared with two different initial velocities, v_1 and v_2, that satisfy $v_2 = 2v_1$. What is the proportion of the periods to complete the cyclotron motion between these two cases? What is the proportion of the radii of the cyclotron orbits between the two cases?

(b) Find a general relation between the classical kinetic energy of the particle and the classical radius of its cyclotron orbit.

(c) Assuming that energy levels are quantized according to $E_n = \hbar\omega_c(n + 1/2)$ where ω_c is the cyclotron frequency, find a heuristic semiclassical relation for the cyclotron radius, R_n, of the n-th orbital, using the result from part (b).

2.2. Electron velocity and conductivity

Consider a classical particle of mass m and charge q moving in two dimensions in the presence of a perpendicular magnetic field, $\mathbf{B} = B\mathbf{e}_z$, and a constant in-plane electric field, $\mathbf{E} = E_x\mathbf{e}_x$.

(a) Write and solve the classical equations of motion for the particle in this case.

(b) Compute the time average of the velocity over one cyclotron period, namely:

$$\langle v_x\rangle = \frac{1}{T}\int_t^{t+T} v_x(t)dt$$

$$\langle v_y\rangle = \frac{1}{T}\int_t^{t+T} v_y(t)dt$$

(c) Define the electrical current density as $j = -en\langle v\rangle$ (here, we choose $q = -e$ for an electron) and compute the classical conductivity matrix, namely, the linear coefficients relating j_x and j_y to E_x.

2.3. Equation of motion

Consider a quantum particle of mass m and charge $q = -e$ moving in two dimensions in the presence of a perpendicular magnetic field, $\mathbf{B} = B\mathbf{e}_z$.

(a) The quantum velocity operators can be defined as follows,

$$\hat{v}_x = \frac{\hat{p}_x + eA_x(\hat{x},\hat{y})}{m}, \qquad \hat{v}_y = \frac{\hat{p}_y + eA_y(\hat{x},\hat{y})}{m}$$

where $\hat{p}_{x,y}$ are the standard momenta that obey the canonical commutation relations with the position operators $\hat{x}$ and $\hat{y}$. Compute the commutator of these velocity operators, $[\hat{v}_x, \hat{v}_y]$.

(b) Define operators for the guiding center,

$$\hat{x}_0 = \hat{x} - \frac{\hat{v}_y}{\omega_c}, \quad \hat{y}_0 = \hat{y} + \frac{\hat{v}_x}{\omega_c}$$

and compute their commutator, $[\hat{x}_0, \hat{y}_0]$.

(c) Derive the Heisenberg equations of motion for the above velocities, for a Hamiltonian that includes the electric field along x,

$$\mathcal{H} = \frac{m}{2}(\hat{v}_x^2 + \hat{v}_y^2) + eE_x\hat{x}.$$

The Heisenberg equation of motion is given by

$$\frac{d\hat{O}}{dt} = \frac{i}{\hbar}[\mathcal{H}, \hat{O}]$$

where $\hat{O}$ is an arbitrary operator.

(d) Computing the equations of motion for the expectation values of the $\hat{v}_x$ and $\hat{v}_y$ operators should return same expressions as in the classical case. However, we know that $\sigma_{xy} = n_e e/B$ classically vs. $\nu e^2/h$ in the quantum case. Show that these two results are in fact equivalent. To this end, express ν via the number of flux quanta.

Problem sheet 2: Landau levels and quantum Hall effect **20.10.2026**

Indicate the completion of the problems 2.1 and 2.2, in percent

	2.1	**2.2**	**2.3**
Uzair Ahmad			
Hamza Akbar			
Sara Arab Badkuba			
Florian Arzt			
Sauman Ashraf			
Nathalie Elisabeth Badt			
Huey Baum			
Damian Bialojahn			
Abhirami Binu			
Ricardo Bouso Ríos			
Syed Muneef Hasan Bukharee			
Calina Burciu			
Manuel Carbonell Diaz			
Anes Cherih			
Om Vedant Chimmalgi			
Attila Dicso			
Egor Dovgoshiy			
Omar Elzeki			
Kaiti Feng			
Jakob Gäbler			
Jamal Ghaith			
Soham Ghatak			
Bruno Hawkes			
Leonard Bendix Heinze			
Anna Louisa Helzel			
Axelle Ievers			
Muhammad Irfan			
Muhammad Zulasheer Kazmi			
Muhammad Zohaib Khalid			

Problem sheet 2: Landau levels and quantum Hall effect **20.10.2026**

Indicate the completion of the problems 2.1 and 2.2, in percent

	2.1	**2.2**	**2.3**
Aleksandre Kobulia			
Tim Koschinsky			
Max Kramer			
Malte Flynn Kretzschmar			
Daniel Lahdo			
Maurice Lindenberg			
Julien Martin			
Alexander Masuda			
Mykhailo Motylets			
Souvik Mukherjee			
Paul Neugebauer			
Daniil Ososkov			
Mikoto Perut			
Mesba Uddin Pias			
Alexandru Plesca			
Hasnat Ahmed Raza			
Guilherme Schewtschik			
Manav Shahaney			
Adil Shahzad			
Mashal Shaikh			
Zhenming Shi			
Faust Siftar			
Valentyn Skalenchuk			
Simona Vasilevska			
Sneha Velmurugan			
Lara Charlène Wollert			
Zijing Yang			
Vanessa Zhilkina			